Ruperto-Carola University Heidelberg
Combined Faculties for Natural Sciences
and for Mathematics

Measurement of Open Charm in 158 AGeV/c Pb-Au Collisions

Dissertation
submitted by
Dipl.-Phys. Wilrid Ludolphs
born in
Königstein im Taunus

February 2006

Referees:

Prof. Dr. Johanna Stachel

Prof. Dr. Franz Eisele

Abstract

Measurement of Open Charm in 158 AGeV/c Pb-Au Collisions

This thesis presents a measurement of an upper limit for the open charm yield in 158 AGeV/c Pb-Au collisions with the CERES spectrometer at 7% centrality. A secondary particle reconstruction scheme, based on the reconstruction of the decay vertex, is developed and tested using the decay $K_S^0 \to \pi^+\pi^-$ as a reference measurement. An integrated K_S^0 rapidity density of $dN/dy = 19.75 \pm 0.23\,(\mathrm{stat}) \pm 1.70\,(\mathrm{syst})$ is measured in the rapidity region $2.0 < y < 2.6$ and compared to results from an alternative analysis of the CERES data and results from the NA49 and NA57 collaborations.

The analysis of the decay $\overline{D}^0 \to K^+\pi^-$ requires careful study of the combinatorial background and resonances contributing to the invariant mass spectrum. An open charm enhancement of more than a factor 22 can be excluded at 98% confidence level. The enhancement is calculated with respect to the expected open charm yield in nucleus-nucleus collisions of $\langle \overline{D}^0 \rangle = 0.21$ per event, obtained by scaling the charm cross-section in proton-proton collisions with the number of binary collisions.

The first part of this thesis is devoted to the development of a hit finding algorithm for the CERES TPC. Furthermore, a detailed description of the calibration of the TPC is presented. A position resolution of $\sigma_{r\Delta\phi} \approx 340$ μm and $\sigma_{\Delta r} \approx 640$ μm is achieved with the new calibration, with a momentum resolution of $\Delta p/p = 5.4\%$ at $p = 5$ GeV/c, the momentum region relevant for the $\overline{D}^0$ meson.

Open Charm Analyse in 158 AGeV/c Pb-Au Kollisionen

Diese Doktorarbeit beinhaltet die Messung einer oberen Grenze für die Produktionsrate von Open Charm Mesonen in 158 AGeV/c Pb-Au Kollisionen mit dem CERES Spektrometer bei einer Zentralität von 7%. Es wird die Entwicklung einer Rekonstruktionsmethode für Sekundärteilchen vorgestellt, die auf der Rekonstruktion des Zerfallsvertex beruht. Die Methode wird mit dem Zerfall $K_S^0 \to \pi^+\pi^-$ als Referenzmessung getestet. Eine integrale Rapiditätsdichte von $dN/dy = 19.75 \pm 0.23\,(\mathrm{stat}) \pm 1.70\,(\mathrm{syst})$ wird für das K_S^0 Meson im Rapiditätsbereich $2.0 < y < 2.6$ gemessen und mit den Ergebnissen einer alternativen CERES Messung und den Messungen der NA49 und NA57 Kollaborationen verglichen.

Eine genaue Untersuchung des kombinatorischen Untergrundes und der Resonanzen, die zusätzlich zu dem invarianten Massenspektrum beitragen, ist für die Analyse des Zerfalls $\overline{D}^0 \to K^+\pi^-$ unerlässlich. Eine vermehrte Open Charm Produktionrate von mehr als einem Faktor 22 kann bei einem Konfidenzniveau von 98% ausgeschlossen werden. Dieser Faktor bezieht sich auf die in Kern-Kern Stößen erwartete Open Charm Produktionsrate von $\langle \overline{D}^0 \rangle = 0.21$ pro Ereignis, die durch Skalierung des Charm Wirkungsquerschnitts in Proton-Proton Stößen mit der Anzahl der binären Kollisionen gegeben ist.

Der erste Teil dieser Doktorabeit befasst sich mit der Entwicklung eines Algorithmus zur Mustererkennung in der CERES TPC. Ferner wird eine detaillierte Beschreibung der Kalibration der TPC präsentiert. Mit der neuen Kalibration wird eine Ortsauflösung von $\sigma_{r\Delta\phi} \approx 340$ μm und $\sigma_{\Delta r} \approx 640$ μm erreicht, sowie eine Impulsauflösung von $\Delta p/p = 5.4\%$ bei einem für das $\overline{D}^0$ Meson relevanten Impuls von $p = 5$ GeV/c.

Contents

List of Figures

List of Tables

Chapter 1

Introduction

Heavy-ion collisions at ultra-relativistic energies offer the possibility to study the behavior of nuclear matter at high density and temperature where one expects the existence of the **Q**uark **G**luon **P**lasma (QGP). In this new state of matter the quarks are no longer confined inside individual hadrons, but they are free to move within the interaction region. At the same time the quarks lose their dynamical mass leading to the restoration of chiral symmetry, an approximate symmetry of QCD in the sector of light quarks.

Many different signatures have been proposed for the formation of a QGP phase. According to the type of final state particles, they can be divided into two main groups: the hadronic probes and the electromagnetic probes. Hadronic probes have the advantage of having large cross-sections. However, they undergo a substantial evolution through strong reinteractions in the period between their formation and the detection. Therefore, the momentum distribution and the final particle composition can be affected by later stages of the heavy-ion collision. Nevertheless, it is still possible to extract valuable information about the initial stage of the collision by studying final state hadrons. Electromagnetic probes have a negligible cross-section for the interaction with hadronic matter. They are a direct tool for the investigation of the first stages of the heavy-ion collision. The drawback of this kind of probe is that the production cross-section is very small, and the detection in a high background environment becomes a difficult task.

CERES is one of the experiments at the CERN **S**uper **P**roton **S**ynchroton (SPS) dedicated to the study of electromagnetic probes. Its main objective is the measurement of low mass electron-positron pairs produced in p-A and A-A collisions. Systematic studies have been done with S-Au in 1992 [1] and the proton-induced reactions p-Be and p-Au in 1993 [2]. An energy scan has been performed during the years 1995 to 2000 using the system Pb-Au [3, 4, 5]. While the p-A collisions could be well explained by a cocktail of the hadronic sources π^0, η, η', ρ, ω and ϕ, the S-Au and Pb-Au spectra showed a significant enhancement in the mass range $0.2 < m_{e^+e^-} < 1.5$ GeV/c^2. A comparison between the p-Au and S-Au data sets is shown in figure 1.1. A dilepton enhancement was also observed by the HELIOS/3 experiment in 200 AGeV/c S-W collisions [6, 7] and by the NA38/NA50 experiment in 158 AGeV/c Pb-Pb collisions [8]. The observation of this excess also led to an enormous theoretical activity. It might be explained by direct thermal radiation of the fireball, dominated by the two-pion annihilation $\pi^+\pi^- \rightarrow \rho \rightarrow e^+e^-$



Figure 1.1: **Low-mass electron pair enhancement in nucleus-nucleus collisions.** *The left figure shows the inclusive e^+e^- mass spectra in 450 GeV/c p-Au collisions. The shape of the spectra is in good agreement with the predicted contributions from hadronic decays. The right figure shows the corresponding spectra for S-Au at 200 AGeV/c. In the mass range between 0.2 and 1.5 GeV/c^2 an enhancement by a factor of* $5.0 \pm 0.7\,(\mathrm{stat}) \pm 2.0\,(\mathrm{syst})$ *[1] is observed with respect to expectations from the hadronic cocktail.*

with an intermediate ρ vector meson. Compared to the lighter vector mesons ω and ϕ the ρ is of particular interest. Because of its short lifetime of 1.3 fm/c it samples the evolution of the fireball as a function of time. Furthermore, due to its link to chiral symmetry restoration [9], it is assumed that the intermediate ρ suffers strong in-medium modifications. The two main theoretical alternatives for this modification predict a shift of the ρ peak to lower masses (Brown-Rho scaling [10]) or a spreading of its width based on calculations of the ρ spectral density within the dense hadronic medium [11]. The upgrade of the CERES experiment in 1998 with a **T**ime **P**rojection **C**hamber (TPC) opened the additional possibility to study hadronic channels. Many interesting topics have been addressed since then [12, 13, 14, 15, 16].

The aforementioned dilepton enhancement in the low and intermediate mass region gave rise to several speculations. One often discussed possibility proposes an abnormally enhanced open charm production in nucleus-nucleus collisions [17]. However, such an increase is not easily justified theoretically [18], and would also be difficult to reconcile with the current understanding of the observed J/Ψ suppression. Another explanation argued that D meson rescattering in hot and dense matter might generate a transverse momentum broadening which can enrich the $\mu^+\mu^-$ phase space covered by the NA50 experiment [19]. But the resulting increase was to small to explain the excess [20]. Furthermore, the role of secondary Drell-Yan processes in hadronic rescatterings has been investigated within the UrQMD transport model [21]. It has been found that in the **i**ntermediate **m**ass **r**egion (IMR) this contribution may constitute up to 30% of the primordial Drell-Yan yield and thus far from explaining the experimental findings. An attempt to explain the HELIOS-3 data used a transport model incorporating dilepton production through secondary hadronic annihilation processes [22, 23]. It was found that the enhancement in S-W collisions could indeed be explained. Unfortunately, the statistics of the HELIOS-3 data for $m_{\mu^+\mu^-} > 2$ GeV is poor. Finally, the significance of thermal dilepton radiation has been investigated [24]. The evaluation of the dilepton spectra in Pb-Pb collisions as measured by NA50 showed that the IMR can be composed of 3 major components: open charm decays, Drell-Yan and thermal radiation. It was shown that the dilepton excess could indeed be explained without invoking any anomalous enhancement in the charm production. Furthermore, the low-mass dilepton spectra from the CERES experiment could be explained in the same framework when medium effects in the low-lying vector mesons are included. Therefore, a consistent picture of dilepton production at the full CERN SPS energy seems to emerge. To test this result against open charm enhancement a direct measurement of the D meson contribution is needed.

The subject of this thesis is the measurement of the open charm yield in 158 AGeV/c Pb-Au collisions. Due to the low production cross-section the measurement is difficult. The expected yield per event for central Pb-Pb collisions at SPS is ~ 0.03 for D^+/D^- and ~ 0.1 for $D^0/\overline{D}^0$ [25, 26]. However, the CERES experiment has the possibility to distinguish between primary and secondary tracks by cutting on the secondary vertex. Moreover, the upgrade of the experiment with a TPC and its extensive calibration provides a reasonable momentum resolution ($\Delta p/p = 5.4\%$ at $p = 5$ GeV/c). Finally, the large data sample of 30 million events taken during the beam time of the year 2000 is enough statistics to make such a measurement tempting.

The thesis is structured as follows. Chapter 2 gives a brief historical overview of the heavy-ion program and elucidates the most relevant features of the Quark Gluon Plasma. Chapter 3 presents the experimental findings of the NA38/NA50 experiment concerning the excess in the intermediate mass muon pair continuum in 158 AGeV/c Pb-Pb collisions. Furthermore, its most discussed sources, open charm enhancement and thermal radiation, are presented. The overall experimental setup of the CERES experiment together with its individual detector components are described in chapter 4. Chapter 5 is devoted to the reconstruction scheme of the raw data. Special emphasis is given to software components developed in the framework of this thesis. Another important contribution during this work was the calibration of the TPC, described in detail in chapter 6. Chapter 7 presents the analysis scheme for the measurement of open charm. It is mainly based on the reconstruction of the secondary vertex using two **S**ilicon **D**rift **C**ounters (SiDC) and the TPC. A cut on the longitudinal distance between the secondary vertex and the primary interaction region allows to separate secondary particles from target tracks and thus makes an almost background free particle reconstruction possible. In chapter 8 the feasibility of the analysis is demonstrated for the more abundant strange particles. The studies are performed on the two body decay $K^0_S \rightarrow \pi^+\pi^-$. The efficiency is tested against an alternative CERES measurement using only the TPC and measurements from the NA49 and NA57 experiments. The same analysis scheme is applied to the reconstruction of open charm in chapter 9. An upper limit on the open charm cross-section is obtained. Finally, chapter 10 contains the conclusions.

Chapter 2

Quark Gluon Plasma

Several experimental discoveries in particle physics revealed the hierarchical nature of matter. The constituents of macroscopic matter were found to be molecules and atoms. The atoms consist of a nucleus surrounded by an electron cloud. The components of the nuclei are the nucleons which in turn are formed of quarks, antiquarks and gluons (partons). To this date no further substructure of quarks was observed. They are regarded as pointlike particles. The properties of the 6 known quark flavors are listed in table 2.1. Isolated quarks have never been detected and thus it is conjectured that they are confined together with other quarks to form hadrons. The gluons are the intermediators of the strong color force between the quarks. The physics of the strong interaction is described by the theory of **Q**uantum **C**hromo**d**ynamics (QCD). It is the SU(3) gauge symmetric part of the Standard Model of particle physics.

In deep-inelastic scattering experiments the individual scattered partons fly away practically freely, dress with a gluon cloud and rapidly form color singlet bound states, the hadrons. However, if the number of partons scattering simultaneously into the same volume element is sufficiently increased, the situation may change radically. A dense medium of partons is formed, where the interactions of quarks, antiquarks and gluons are screened such that the formation of bound states is inhibited. This new state of matter is called Quark Gluon Plasma.

The ultimate aim of ultra-relativistic heavy-ion collisions is to prove the existence of this new state of matter and to investigate its particular properties. The topic is also relevant to other fields, like cosmology and astrophysics. Firstly, the transition from hadronic matter to a QGP may happen in the inner core of neutron stars, where mass densities are likely to exceed 10^{15} g/cm^3 (about four times the central density of nuclei) while surface temperatures are as low as 10^5 K or less [11]. Secondly, it is believed that the inverse transition (hadro-synthesis) had to occur a few tens of microseconds after the Big Bang. The following sections will give a brief historical overview over the heavy-ion program and elucidate the most interesting features of the phase transition.

quark flavor	charge Q/e	current mass m_q [MeV/c^2]	dynamical mass M_q [MeV/c^2]
d (down)	$-1/3$	9.9 ± 1.1	310
u (up)	$+2/3$	5.6 ± 1.1	310
s (strange)	$-1/3$	199 ± 33	500
c (charm)	$+2/3$	1350 ± 50	1800
b (bottom)	$-1/3$	$\simeq 5000$	4500
t (top)	$+2/3$	> 90000	> 176000

Table 2.1: **Quark properties.** *Quarks are fermions (spin 1/2) and are characterized by the flavor degree of freedom d, u, s, c, b or t. The current quark masses listed in the table were taken from reference [27], the dynamical masses from [28]. A current mass is the mass of a quark in the absence of confinement. When the quark is confined in a hadron, it may acquire an effective mass which includes the effect of the zero-point energy of the quark in the confining potential. This so called dynamical mass is typically a few hundred MeV in magnitude.*

2.1 Historical Overview

The heavy-ion program started with fixed target experiments in 1986 at the **A**lternating **G**radient **S**ynchroton (AGS) of the **B**rookhaven **N**ational **L**aboratory (BNL) and the **S**uper **P**roton **S**ynchroton (SPS) of the **C**entre **E**uropéen pour la **R**echerche **N**ucléaire (CERN). The center of mass energy reached with these two facilities is $\sqrt{s} = 5$ AGeV and $\sqrt{s} = 17$ AGeV, respectively. Data was taken using a beam of relatively light ions of 14.6 AGeV ^{16}O and ^{28}Si at the AGS and 60-200 AGeV ^{16}O and ^{32}S beams at the SPS. It was observed that the colliding nuclear matter loses a substantial fraction of its energy in the collision process [29]. This proved the possibility to create hot and dense matter in heavy-ion collisions.

The program continued with heavier nuclei, 10.8 AGeV ^{197}Au at the AGS and 40, 80 and 158 AGeV ^{208}Pb at the SPS. The high energy Pb data provided circumstantial evidences that a new state of matter had been produced [30], and the official announcement from CERN soon followed in the year 2000 [31].

The QGP formed at SPS energies is not expected to be net-baryon-free. The net-baryon content in the QGP is small, if the separation between the beam and the target rapidity stays far away from the central rapidity region. For nucleus-nucleus collisions at an energy of $\sqrt{s} = 100$ AGeV, the separation between the projectile and the target rapidity is 10.7 units. On the other hand, the average rapidity loss of the baryons in the central collision of Au on Au is roughly 2 to 4 units [27]. Thus, for those high energy collisions a net-baryon-free Quark Gluon Plasma would be expected. This triggered the set of experiments, which began to run in the year 2000 at the **R**elativistic **H**eavy **I**on **C**ollider (RHIC), also located at BNL. These were collider experiments, able to reach even higher energies of about $\sqrt{s} = 200$ AGeV necessary to produce a net-baryon-free Quark Gluon Plasma.

An even clearer result will be obtained with a new collider being built at CERN, the **L**arge **H**adron **C**ollider (LHC). This machine will be capable to reach energies of about $\sqrt{s} = 5.5$ ATeV. At this highest energy all parameters relevant for the formation of the



Figure 2.1: **Phase diagram of matter.** *Lattice QCD calculations predict a deconfinement phase transition, and an accompanying chiral phase transition, both at high temperature and/or high density. The expectation for the phase boundary [35], based on calculation for $\mu_B = 0$, is indicated by the hatched region. The chemical potentials μ and temperatures T resulting from thermal analyses place the chemical freeze-out (black line) very close to the phase boundary between plasma and hadrons.*

QGP, such as energy density, size and lifetime of the system and relaxation time, will be more favorable. The experiment dedicated to heavy-ion collisions at LHC will be ALICE (**A** **L**arge **I**on **C**ollider **E**xperiment) [32]. First runs are planned for the year 2007.

2.2 Phase Diagram of Nuclear Matter

Recent theoretical predictions estimate that the phase transition between the confined and the deconfined phases of QCD occurs at a critical energy density of $\epsilon_c = 0.70 \pm 0.35$ GeV/fm^3 [33]. This energy density can be reached by changing one or both of the two essential thermodynamical quantities: the temperature T and the density ρ. The QCD phase diagram in figure 2.1 can thus map out regions in the plane of temperature T and baryochemical potential μ_B [34], with the latter specifying the mean



Figure 2.2: **Space-Time Evolution of a Heavy-Ion Collision.** *The invariant hyperbola in the space-time diagram allocate the individual stages passed during a heavy-ion collision.*

baryon number density ρ_B (baryon minus antibaryons).

If ordinary nuclear matter is compressed to the extent that nucleons overlap, a cold QGP is formed. It is believed that this situation may take place inside neutron stars [36]. At $T = 0$ the critical density for the transition can be estimated to be $\rho_c \simeq 0.72/\mathrm{fm}^3$ [27]. ρ_c is about 5 times the nucleon number density $\rho_B = 0.14/\mathrm{fm}^3$ for normal nuclear matter at equilibrium. Experimentally, the QGP can be accessed by a combination of heating and compression, colliding heavy nuclei with velocities close to the speed of light. The energy brought into the system is dissipated for the creation of quark-antiquark pairs and gluons. Assuming equilibration, the system becomes deconfined at sufficiently high density and temperature to form a hot QGP.

The space-time scenario of a high-energy nucleus-nucleus collision was suggested by Bjorken [37] as illustrated in figure 2.2. The trajectories of the projectile and the target nucleus are drawn as thick lines in the diagram. The nuclei are Lorentz contracted in longitudinal direction and therefore represented by two thin disks. During the collision process the baryons lose a substantial fraction of their energy, which is deposited in the vicinity of the center of mass. After the collision the slowed-down baryons can still have enough momentum to recede from each other. The energy deposition is approximately additive in nature and might be sufficient to create a system of quasi-free quarks and gluons in the central rapidity region. After a short moment of incomplete thermal equilibration (pre-equilibration), the plasma may reach local equilibration (QGP) at the formation time

$\tau_0 \approx 1$ fm/c [37]. The subsequent evolution of the fireball can be described in terms of the laws of hydrodynamics. The pressure inside the hot and dense medium leads the system to expand. Its temperature drops and the plasma starts to hadronize (mixed phase). The hadrons will stream out of the collision region when the temperature falls below the temperature of chemical freeze-out. At this stage all inelastic interactions stop and the yield of each hadron species is fixed. As the temperature further decreases to reach the point of thermal freeze-out, also the elastic collisions cease.

The relative abundance of produced hadrons in heavy-ion collisions can be compared with expectations for a statistical ensemble [34]. This allows two parameters to be adjusted to the data: the temperature T and the baryochemical potential μ_B. At present, data exists for a variety of energy regimes, accessible by different accelerators (SIS, AGS, SPS and RHIC). This allowed the determination of hadrochemical freeze-out points, shown in figure 2.1 as red circles. The empirical curve connecting the freeze-out points corresponds to a constant energy per hadron of 1 GeV [38]. The information about the thermal freeze-out can be extracted from the analysis of momentum spectra and is also shown in the figure.

2.3 Chiral Symmetry Restoration

A fundamental symmetry of QCD is chiral symmetry. It is broken in the vacuum, but lattice QCD predicts that chiral symmetry is restored at the same temperature as the deconfinement of the quarks and gluons occurs. An important goal in heavy-ion physics is therefore to look for evidences for this fundamental prediction of the Standard Model. This issue will briefly be elucidated in the present section. A comprehensive introduction to this topic can also be found in [39].

Chiral symmetry involves the handedness of particles, i.e. the information of the spin orientation versus its direction of motion. The ability to define the handedness in an absolute way depends on weather or not a particle has mass, since a particle with mass moves with less than the speed of light. For right-handed particles, the spin is pointing in the same direction as its velocity. But for an observer moving faster than the particle, it would appear to move backwards, while its spin would still remain unchanged. For that observer the particle would seem to be left-handed. In contrast, a massless particle moves at the speed of light in all frames, and all observers would agree on its handedness.

As indicated in table 2.1 the masses of the quarks are not fixed constants but are rather generated by interactions with other particles. The current mass m_q can be associated with the weak interaction. The dynamical mass M_q arises from the strong interaction. It is believed that the current masses are generated by the coupling to the still undiscovered Higgs field. The Higgs field has a finite average value in the ground state, generally known as vacuum. Particles interact with the vacuum Higgs field, which modifies their properties and gives rise to their masses. In the limit $m_q \to 0$ the QCD Lagrangian for light quarks (u,d, and s) reveals SU(3) flavor symmetry independently for left-handed and right-handed quarks, i.e. it has chiral symmetry $\mathrm{SU(3)_L} \times \mathrm{SU(3)_R}$. In this limit the helicity becomes a good quantum number. Since the actual current masses of the light quarks



Figure 2.3: **Energy dependence of a charge singlet pair.** *Due to quantum fluctuations the vacuum is filled with pairs of charged particles in a singlet state. Charge means electric charge in the case of QED (***Q***uantum* **E***lectro***d***ynamics) and color charge in the case of QCD (***Q***uantum* **C***hromo***d***ynamics). If a e^+e^- pair is created from the QED vacuum, it will annihilate within the time scale $1/E_{pair}$ given by the uncertainty relation. These are virtual pairs. In contrast to this scenario, the total energy in QCD exhibits a minimum at negative energy. The empty vacuum at $E = 0$ becomes unstable and the pairs become real.*

are small compared to the typical few hundreds of MeV for strongly interacting particles, they can to a good approximation be regarded as massless and QCD as approximately invariant under the chiral version of isospin symmetry.

As is schematically illustrated in figure 2.3, the total pair energy in QCD has a minimum at some distance $r_0 \sim 1$ fm. Moreover the value of this minimum is negative. As a consequence, an empty ($E = 0$) vacuum becomes unstable because there exists a configuration with lower energy. Pairs of color charge are created and stay there forever. The QCD vacuum is filled with real color charge pairs. Because the gauge field (gluons) is a vector field, due to the helicity conservation for $m_q \to 0$ a left-handed quark can only couple with a left-handed antiquark and vice versa. This means that the $q\bar{q}$ pairs in the QCD vacuum have to be in the singlet state not only in color but also in spin. But already this means that chiral symmetry is broken in the vacuum, since there are $q\bar{q}$ pairs in a scalar state, i.e. containing q and $\bar{q}$ of opposite helicities, which in massless limit do not interact. A right-handed and massless test quark put into such a vacuum can only annihilate on a right-handed antiquark thus liberating a left-handed quark [40]. For an observer at some distance this will look like the test quark, being in a vacuum, changes spontaneously its helicity. Therefore, it cannot move with the speed of light, and hence it had to acquire some dynamical mass. It is these interactions with the asymmetric vacuum that hide the approximate chiral symmetry of QCD.

If the temperature and therefore the kinetic energy is raised, at some critical value T_c of the order of the pion mass the energy stored in the strong field is overcome. At this stage the minimum of the total pair energy will become positive, and hence the real $q\bar{q}$ pairs would disappear from the vacuum. Above that temperature chiral symmetry will be restored and quarks will retain their zero mass in the chiral limit.

2.4 Equation of State

The equation of state describes the temperature dependence of the energy density ϵ and the pressure p. Obviously it is of great interest to estimate the hadron-quark transition temperature. The simplest picture is perhaps assuming the hadronic phase to be an ideal non-interacting **H**adron **G**as (HG) of massless pions, and the quark phase to be an ideal gas (QGP) of massless gluons and two-flavor quarks [41]. The energy density ϵ and the pressure p of an ideal gas are given by:

$$\epsilon_{ideal} = \frac{g}{30}\pi^2 T^4, \tag{2.1}$$

$$p_{ideal} = \frac{\epsilon_{ideal}}{3} = \frac{g}{90}\pi^2 T^4. \tag{2.2}$$

The degeneracy factor g is defined by the boson degrees of freedom n_b and the fermion degrees of freedom n_f:

$$g = n_b + \left(1 - \frac{1}{2^3}\right) n_f. \tag{2.3}$$

With the boson degrees of freedom $n_b = 3$ for the three isospin pion states, and no fermions, the pressure of the ideal Hadron Gas would be:

$$p_{HG} = \frac{1}{30}\pi^2 T^4 \simeq \frac{1}{3} T^4. \tag{2.4}$$

In the QGP case, the boson degrees of freedom add up to $n_b = 16$ (8 color states $\times$ 2 spin states for gluons). The fermion degrees of freedom add up to $n_f = 24$ (2 flavors $\times$ 3 colors $\times$ 2 spins $\times$ 2 for quark-antiquark). The external pressure can be estimated using the bag model for hadrons [42]. It is equal to the bag constant B. As a result, the pressure for the QGP can be written as:

$$p_{QGP} = \frac{37}{90}\pi^2 T^4 - B \simeq 4T^4 - B. \tag{2.5}$$

For low temperatures, equation (2.4) yields the larger pressure. By increasing the temperature, the two pressures first become equal at a critical temperature T_c and thereafter the QGP pressure dominates. According to Gibbs criterion, the phase with the largest pressure is the stable one. This means that at low temperature the system will be in a HG phase and for high temperatures in the QGP phase. Assuming a bag constant of $B^{1/4} \simeq 200$ MeV [27], the transition will occur at a temperature:

$$T_c = \left(\frac{90B}{37\pi^2}\right)^{\frac{1}{4}} \approx 140MeV \tag{2.6}$$

More detailed theoretical calculations are done in lattice QCD, a nonperturbative treatment of quantum chromodynamics formulated in a discrete lattice of space-time coordinates [43]. They have been performed at zero net-baryon charge. This has a pure technical reason, as some expressions that are in lattice Monte Carlo formulation interpreted as probabilities, are no longer positive definite at non-zero baryon densities.



Figure 2.4: **Energy density and pressure as a function of temperature in lattice QCD.** *The calculations have been performed by the Bielefeld group [33] using 0, 2 and 3 light quarks as well as two light quarks and a heavier (strange) quark. At high temperature it is expected that ϵ/T^4 and p/T^4 will asymptotically approach the free gas limit for a gas of gluons and n_f quarks (equation 2.1 and 2.2). This is indicated by the arrows.*

Figure 2.4 shows a calculation for the energy density and the pressure. The rapid change in energy density within a narrow temperature interval indicates the change from confined hadrons to a QGP. However, the phase transition is found to be of the first or of the second order, or even a smooth cross-over, depending on the number of quark flavors and their masses. The critical temperature T_c obtained at zero baryon density is about 260 MeV for pure gauge theory, i.e. for systems consisting of gluons only. For theories with dynamical quarks T_c varies between 140-170 MeV.

Chapter 3

Dimuon Excess in the Intermediate Mass Region

Dileptons have the interesting property of not suffering strong interactions with the surrounding hadronic medium. Therefore, they are considered as ideal probes of the early stage of heavy-ion collisions, where the Quark Gluon Plasma formation is expected. The dilepton mass spectra can basically be divided into three regions [22]. The region below the ϕ meson (~ 1 GeV/c^2) is addressed as the low mass region. It is dominated by hadronic interactions and hadronic decays. In the intermediate mass region from about 1 to 2.5 GeV/c^2, the contribution from the thermalized QGP might be seen [44]. And finally, in the high mass region the J/ψ suppression has been a subject of great interest, since it was proposed as a signal of the deconfinement phase transition [45].

The most prominent observation in the low and intermediate mass region has been the dilepton enhancement seen by the experiments HELIOS/3, CERES and NA38/NA50. The NA50 experiment has proposed an enhanced open charm production in central Pb-Pb collisions by a factor of 3 [17]. Such a large enhancement is, however, difficult to explain theoretically. Another interpretation of the data is that the increase of the IMR dimuons is not from charm but is due to thermal dileptons [24]. This chapter summarizes the experimental findings and gives a brief confrontation of two possible explanations: open charm enhancement and thermal dilepton radiation.

3.1 Experimental Evidences

The NA38/NA50 experiment studied muon pair production in p-A, S-U and Pb-Pb collisions at the CERN SPS [8, 46]. The dominant contribution to the dimuon spectrum in the IMR is the combinatorial background due to the decays of π and K mesons. After its subtraction the invariant mass and transverse momentum spectra in p-A collisions are well described by a superposition of dimuons coming from leptonic and semi-leptonic charm decays and from Drell-Yan processes:

$$\frac{dN^{\pm}}{dm} - R\frac{dN^{bg}}{dm} = n^{J/\psi}\frac{dN^{J/\psi}}{dm} + n^{\psi'}\frac{dN^{\psi'}}{dm} + n^{D\overline{D}}\frac{dN^{D\overline{D}}}{dm} + n^{DY}\frac{dN^{DY}}{dm} \tag{3.1}$$



Figure 3.1: **Charm hadroproduction cross-section.** *The values refer to forward production ($x_F > 0$). The dotted curve shows the $\sqrt{s}-$dependence of $\sigma_{c\bar{c}}^{pp}$ obtained with PYTHIA and fitted to the measurements. The red square is the indirect measurement from the NA38/NA50 collaboration [17].*

In this equation R is a correction factor to account for charge correlation effects. The shape of the dimuon sources in the right part of equation (3.1) is evaluated by means of a Monte Carlo simulation. By fixing the shape of the dimuon sources, the p-A spectra are fitted leaving the yields n^i ($i = DY$, $D\overline{D}$, J/ψ, ψ') as free parameters.

The obtained total number of muon pairs from $D\overline{D}$ decays is translated into an open charm production cross-section $\sigma^{pA}_{D\overline{D}\to\mu\mu}$ for p-A collisions. To obtain the corresponding value for p-p collisions at 450 GeV/c a linear dependence on the mass number A is assumed:

$$\sigma_{c\bar{c}}^{pp} = \left(\frac{\sigma_{c\bar{c}}^{pp}}{\sigma^{pp}_{D\overline{D}\to\mu\mu}}\right)^{MC} \cdot \frac{\sigma^{pA,Data}_{D\overline{D}\to\mu\mu}}{A} = 36.2 \pm 9.1\ \mu\text{b}. \tag{3.2}$$

The Monte Carlo scaling ratio was obtained from a simulation with the event generator PYTHIA [47]. The NA38/NA50 compared their indirect measurement for the p-p open charm production cross-section to direct measurements from other experiments, as shown in figure 3.1. For this purpose the cross-section value was divided by a factor of 2 to consider only the $x_F > 0$ hemisphere. The agreement is satisfactory. It is possible to obtain the $\sigma_{c\bar{c}}^{pp}$ cross-section for different energies from a PYTHIA calculation fitted to the data.

The expected cross-section for A-A collisions $\sigma^{AA}_{c\bar{c}\to\mu\mu}$ is obtained by using a linear extrapolation of the p-A sources with the product $A_p \times A_t$ of the mass number of the projectile and target nuclei. This enables to calculate the expected number of $D\overline{D}$ dimuons in the centrality integrated S-U and Pb-Pb spectra. The Drell-Yan events are determined by a fit to the high mass region of the spectrum.