

Quark-Gluon Plasma Physics

7. Correlations and Fluctuations


**Prof. Dr. Peter Braun-Munzinger
Prof. Dr. Klaus Reygers
Prof. Dr. Johanna Stachel
Heidelberg University
SS 2023**

Correlations of identical bosons due to quantum interference

stochastic emission from extended source

consider 2 identical bosons (photons, pions, ...)

2 detectors in locations r_1, r_2 observe identical bosons of momenta p_1 and p_2



cannot distinguish solid and dashed paths because of identical particles

→ for plane waves, the probability amplitude for detection of the pair is

$$A_{12} = \frac{1}{\sqrt{2}} [e^{ip_1(r_1-x)} e^{ip_2(r_2-y)} + e^{ip_1(r_1-y)} e^{ip_2(r_2-x)}]$$

with 4-vectors p, r, x, y (to be general for nonstatic source)

square of amplitude: intensity → “intensity interferometry”

probability $A_{12}^2 \propto (1 + \cos(\Delta k \cdot \Delta x))$

is enhanced for momenta close to each other due to Bose-Einstein statistics

Hanbury-Brown Twiss effect


technique of intensity interferometry developed by Hanbury-Brown and Twiss in astrophysics as a means to determine size of distant objects

R. Hanbury-Brown and R.Q. Twiss, Phil. Mag. 45 (1954) 663

radiowaves to determine size of galaxies (Cygnus, Cassiopeia)

- this could be done with classical Michelson (amplitude) interferometry and Nature 178 (1956) 1046


visible light to determine the size of stars (Sirius)



correlated signal varies on characteristic length scale: $D = \lambda \frac{R_{\text{star}}}{L}$
varying separation of detectors one learns the apparent angle between the sources

➡ with detector separation of the order of meters, star size of order 10^6 km 1 light year away can be determined by intensity interferometry with visible light



Angular diameter of Sirius from HBT Correlations



A TEST OF A NEW TYPE OF STELLAR INTERFEROMETER ON SIRIUS

By R. HANBURY BROWN
 Jodrell Bank Experimental Station, University of Manchester
 AND
 DR. R. Q. TWISS
 Services Electronics Research Laboratory, Baldock
 Nature, Nov. 10, 1956, Vol. 178

intensity interferometry less sensitive to vibrations and atmospheric fluctuations than classical Michelson interferometer



➡ Angular diameter R/L of Sirius from intensity interferometry: $3.1 \cdot 10^{-8}$ rad


HBT interference in particle physics

when phase space volume smaller than $\Delta p_x \Delta x \approx \hbar$ is considered, chaotic system of identical noninteracting particles exhibits quantum fluctuations following Bose-Einstein (or Fermi-Dirac) statistics

first observation in particle physics: **pions with small relative momenta**

G. Goldhaber, S. Goldhaber, W.Y. Lee, A. Pais, Phys. Rev. 120 (1960) 300

pion correlations after a pbarp annihilation



angle between 2 pions in the p pbar cm system

look at angular opening between like- and unlike-sign pions in events with 4 charged pions at $\sqrt{s} = 2.1 \text{ GeV}$

radius r of volume of strong interaction in units of Compton wavelengths of pion $\frac{\hbar c}{mc^2} = 1.4 \text{ fm}$

found to be 0.5 - 0.75
i.e. $r = 0.7\text{-}1.0 \text{ fm}$

HBT interferometry in nuclear and particle physics

after discovery of GGLP systematically used as tool to determine source size in particular in collisions of high energy nuclei

useful references:

G.I. Kopylov, M.I. Podgoretsky, Sov. J. Nucl. Phys. 15 (1972) 219 and ibid 18 (1974) 336 and PLB50 (1974) 472

theory of interference of identical pions, lifetime of source

E. Shuryak, Phys. Lett. B44 (1973) 387

introduces time, i.e. duration of emission


excellent review by: G. Baym, Acta Phys. Polon. B29 (1998) 1839

2-particle correlation function

typical particle physics experiment: vary momenta at fixed location of detector

$$C_2(\vec{p}_1 - \vec{p}_2) = \frac{d^6 N / d\vec{p}_1 d\vec{p}_2}{d^3 N / d\vec{p}_1 \cdot d^3 N / d\vec{p}_2} = 1 + f(\vec{p}_1 - \vec{p}_2)$$

get uncorrelated denominator typically from event mixing



in heavy ion collisions typical dimensions 1-10 fm,
—▶ leads to interference at momentum differences of 20-200 MeV/c

for a stochastic source we get a 2-particle correlation function

$$C_2(q) = P_{12} = \int d^4x d^4y |A_{12}|^2 \rho(x)\rho(y)$$

with a 4-momentum difference $q \equiv p_1 - p_2$

and momentum independent source functions $\rho(x), \rho(y)$

transformation to relative coordinates and integration leads to

$$C_2(q) = 1 + |\tilde{\rho}(q)|^2$$

$$\tilde{\rho}(q) = \int d^4x \rho(x) e^{-iqx}$$

Fourier transform of space-time density distribution

example: Gaussian space-time density distribution $\rho(x)$

$$\rho(x) = c \exp\left(-\frac{|\vec{x}|^2}{2r_0^2} - \frac{t^2}{2\tau_0^2}\right)$$

$$\longrightarrow \tilde{\rho}(q) = c' \exp\left(-\frac{|\vec{q}|^2 r_0^2}{2} - \frac{q_0^2 \tau_0^2}{2}\right)$$

and $C_2(q) = 1 + \exp(-r_0^2 |\vec{q}|^2 - \tau_0^2 q_0^2)$ with norm $\tilde{\rho}(0) = 1$

note that the 3d rms radius is $r_{rms} = \sqrt{3}r_0$

Coupling of spatial information and time

time and space information are coupled, i.e. q_0 and $q(\text{vector})$ are not independent in the pair rest frame $q^* = (0, \vec{q}^*) = (0, 2\vec{k}^*)$ with $\vec{k}^* = \vec{p}_1^* = -\vec{p}_2^*$

$$\text{(in general } \vec{k}^* = \frac{m_2 \vec{p}_1^* - m_1 \vec{p}_2^*}{m_1 + m_2} \text{)}$$

if \tilde{v} is the velocity of the pair in the overall cm frame, $\gamma^2 = \frac{1}{1 - v^2/c^2}$

and $\vec{k}_T^* = \vec{q}_T/2$, $k_L^* = q_L/2\gamma$ (here T and L refer to the direction of the pair)

and $\vec{v} \cdot \vec{q} = v q_L$ but $\frac{\vec{v}}{c} = \frac{\vec{p}_1 + \vec{p}_2}{E_1 + E_2}$

$$\rightarrow \vec{v} \cdot \vec{q} = E_1 - E_2 = q_0$$

$$\rightarrow C_2(q) = 1 + \exp(-4r_0^2 k_T^{*2} - 4\rho^2 \gamma^2 k_L^{*2})$$

where $\rho^2 = r_0^2 + (v\tau_0)^2$

in direction of the pair, spatial and temporal information cannot be distinguished for identical particles

Coulomb correction

in addition to quantum correlations, 2 identical pions also experience correlations due to electromagnetic interaction, i.e. Coulomb repulsion

for 2 pions from a point source this problem was already solved for beta-decay (Gamov function)

$$\psi_C(0) = \left(\frac{2\pi\eta}{e^{2\pi\eta} - 1} \right)^{1/2} \quad \text{with} \quad \eta = \frac{zz'e^2}{v_{\text{rel}}}$$


was used also in early days of HBT, but approximation is too crude

(P. Braun-Munzinger, G. Baym, Nucl. Phys. A610 (1996) 286c)
look at $\pi^+\pi^-$ correlation to see Coulomb effect

need to integrate over source of proper size

examples $R = 1, 5, 9, 18$ fm

a recursive problem, since source size from HBT is determined after Coulomb correction
– typically very few iteration steps



3-dimensional correlations functions in pp and heavy ion collisions

go from momentum difference in cartesian coordinates to so-called Bertsch-Pratt variables (G. Bertsch, Phys. Rev. C37 (1988) 1896)

define z-axis as beam axis

$$q_{long} = \vec{q} \cdot \hat{e}_z = q_z$$

and in transverse direction distinguish direction parallel to pair momentum


$$\hat{e}_{out} = \frac{\vec{p}_{t1} + \vec{p}_{t2}}{|\vec{p}_{t1} + \vec{p}_{t2}|}$$

$$q_{out} = \vec{q} \cdot \hat{e}_{out}$$

and perpendicular to it


$$\hat{e}_{side} = \hat{e}_{out} \times \hat{e}_z$$

$$q_{side} = \vec{q} \cdot \hat{e}_{side}$$



Example: raw measured pp correlation function

CERES/NA45 at CERN SPS - central 158 A GeV/c PbAu collisions



characteristic width order 30 MeV/c

fit function of form $C_2 = 1 + \lambda \exp(-R_{long}^2 q_{long}^2 - R_{side}^2 q_{side}^2 - R_{out}^2 q_{out}^2)$
to the exp. correlation function


with correlation strength $\lambda = 1$ for chaotic source and
0 for fully coherent source

Dynamical source with space-momentum correlations


collective expansion introduces space-momentum correlations leading to an apparent reduction in source radius

simple example:

source with $T=0$ emitting particles radially with a Hubble-like velocity profile



the apparent source from 2-particle correlations looks pointlike!
only particles emitted from the same space point have the same momentum, i.e. a small momentum difference



Dynamical source with space-momentum correlations

effect of hydrodynamic expansion on radius parameters extracted from 2-pion correlations

S. Pratt, Phys. Rev. Lett. 53 (1984) 1219

A.N. Makhlin, Y.M. Sinyukov, Z. Physik C39 (1988) 69

Y.M. Sinyukov, Nucl. Phys. A498 (1989) 151c

picture: thermal source that expands hydrodynamically with velocity $v(r)$ and corresponding $\beta(r)$ and $\gamma(r)$

look at 2-pion correlation function for different slices in pair transverse momentum


$$\vec{k} = \frac{1}{2}(\vec{k}_1 + \vec{k}_2) \quad \text{or also} \quad k_t = \frac{1}{2}(k_{t,1} + k_{t,2})$$

find that $R(k)$ decreases with k , as ratio of energy of collective expansion over thermal energy increases

$$R(k) = R[(y \tanh y)^{-1} - \sinh^{-2} y]$$

$$\text{with} \quad y = k\gamma\beta/T$$


faster pions are more likely to be emitted near the point on spherical shell expanding with velocity β in direction of k (S.Pratt)



First successful hydrodynamic description of HBT data

data: NA49 at SPS, 200 GeV/c Pb+Pb
thesis H. Appelshäuser
Eur. Phys. J. C2 (1998) 661

$$R_{\perp} = \sigma_x \left(1 + \frac{m_t \beta^2}{T \cosh(y_{YPK} - y_{\pi\pi})} \right)^{-1/2}$$




fit function U.Heinz et al.
expanding source with
expansion velocity b ,
temperature T and
pair transverse mass

$$m_t^2 = m_\pi^2 + k_t^2$$



consistent description w.
 $\beta = 0.5$, $T = 120 \text{ MeV}$
 $\sigma_x = 8.2 \text{ fm}$

Velocity of source emitting pions relative to pair velocity



consistent with boost invariant expansion over ± 2 units of y (source rapidity equal pair rapidity)
central fireball with Bjorken expansion 4 units long

R_{long} - Longitudinal Expansion of Fireball

due to finite T , pions are correlated over an interval in rapidity and due to expansion this grows linearly with duration of expansion
 R_{long} is longitudinal correlation length


$$R_{long} \approx \tau \underbrace{\sqrt{T_f/m_t}}_{\text{thermal velocity}} \quad \text{Y.Sinyukov}$$

duration of expansion (lifetime) τ of the system can be estimated from the transverse momentum dependence of R_{long}

$$\tau = 6 - 8 \text{ fm}/c$$

for $T_f = 120 - 160 \text{ MeV}$


CERES Pb-Au *Nucl. Phys. A714 (2003) 124*



Hubble plot of nuclear fireball

Freeze-out volume vs. beam energy

estimate freeze-out volume V_f : $V_f = (2\pi)^{3/2} R_{side}^2 R_{long}$




note: this is volume of 0.88 units of rapidity

- ◆ **surprise:** non-monotonic behaviour
- ◆ **minimum** between AGS and SPS
- ◆ explained by falling nucleon density and increasing pion density together with $\sigma_{\pi n} \gg \sigma_{\pi\pi}$

Densities at chemical and thermal freeze-out


HBT gives density at thermal freeze-out



volume appears only to grow 30 % between chemical and thermal freeze-out
 get from the 2 densities, the velocity and final radii: isentropic expansion for 0.9-2.3 fm
 $T_f = 158 - 132$ MeV, rate of cooling near T_c : (13 ± 1) % per fm/c

ALICE PbPb collisions at the LHC at $\sqrt{s_{nn}} = 2.76$ TeV

ALICE, Phys. Lett. B696 (2011) 328



HBT radii larger at LHC than at RHIC
larger radial flow


k_T dependence shows shape
typical for hydrodynamically
expanding source
rather similar to RHIC data
but radii are larger
 R_{out}/R_{side} dropping below 1
reproduced reasonably well by
Krakow and Kiev hydro models

hydro models

HBT radii much bigger than 1-d charge
radii of colliding nuclei

Updated energy dependence of HBT radii

Phys. Lett. B696(2011)328




$$\langle kt \rangle = 0.3 \text{ GeV}/c$$

- significant extension of reach of world data by ALICE at higher energies radii grow as cube root of multiplicity indicating freeze-out at constant density
- hydro models reproduce growth reasonably well

freeze-out volume and duration of expansion


coherence volume $V = (2\pi)^{3/2} R_{\text{side}}^2 R_{\text{long}}$

from R_{long} : expansion at LHC 10 fm/c



huge growth at LHC


$$R_{\text{long}} = \tau_f \sqrt{T/m_t}$$



ALICE, Phys. Lett. B696(2011)328

Volume in pp and heavy ion collisions different

ALICE Phys. Rev. D84 (2011) 112004



PbPb coll.scale with cube root of multiplicity i.e. freeze-out at constant density for different centralities and energies
pp falls on different curves, i.e. at a given multiplicity pp physics different